



Rectal Ointment

Common Use: This is a topical ointment used for rectal conditions such as anal fissures and hemorrhoids.

Directions: Wash hands before and after use. When applying the medication, be sure to use a finger cot, glove, or Q-tip and apply into anal canal or external area. This insures that the medication is being absorbed only where needed.

Precaution: Apply medication exactly as directed by your physician. Do not use differently without your physicians authorization. Possible side effects from active ingredients include:

Hydrocortisone - Stinging, burning, irritation, dryness, or redness.

Lidocaine - Mild bruising, redness, itching, swelling, mild dizziness, nausea, and numbness.

Nitroglycerin - Dizziness, lightheadedness, syncope, vertigo, drowsiness, and headache.

Diltiazem - Dizziness, lightheadedness, weakness, tired feeling, nausea, upset stomach, flushing (warmth, redness, or tingly feeling), sore throat, cough, stuffy nose, and headache.

Nifedipine - Dizziness, flushing, weakness, swelling ankles/feet, constipation, and headache.



Rectal Ointment

Common Use: This is a topical ointment used for rectal conditions such as anal fissures and hemorrhoids.

Directions: Wash hands before and after use. When applying the medication, be sure to use a finger cot, glove, or Q-tip and apply into anal canal or external area. This insures that the medication is being absorbed only where needed.

Precaution: Apply medication exactly as directed by your physician. Do not use differently without your physicians authorization. Possible side effects from active ingredients include:

Hydrocortisone - Stinging, burning, irritation, dryness, or redness.

Lidocaine - Mild bruising, redness, itching, swelling, mild dizziness, nausea, and numbness.

Nitroglycerin - Dizziness, lightheadedness, syncope, vertigo, drowsiness, and headache.

Diltiazem - Dizziness, lightheadedness, weakness, tired feeling, nausea, upset stomach, flushing (warmth, redness, or tingly feeling), sore throat, cough, stuffy nose, and headache.

Nifedipine - Dizziness, flushing, weakness, swelling ankles/feet, constipation, and headache.



Rectal Ointment

Common Use: This is a topical ointment used for rectal conditions such as anal fissures and hemorrhoids.

Directions: Wash hands before and after use. When applying the medication, be sure to use a finger cot, glove, or Q-tip and apply into anal canal or external area. This insures that the medication is being absorbed only where needed.

Precaution: Apply medication exactly as directed by your physician. Do not use differently without your physicians authorization. Possible side effects from active ingredients include:

Hydrocortisone - Stinging, burning, irritation, dryness, or redness.

Lidocaine - Mild bruising, redness, itching, swelling, mild dizziness, nausea, and numbness.

Nitroglycerin - Dizziness, lightheadedness, syncope, vertigo, drowsiness, and headache.

Diltiazem - Dizziness, lightheadedness, weakness, tired feeling, nausea, upset stomach, flushing (warmth, redness, or tingly feeling), sore throat, cough, stuffy nose, and headache.

Nifedipine - Dizziness, flushing, weakness, swelling ankles/feet, constipation, and headache.